

**MARK SCHEME for the May/June 2009 question paper
for the guidance of teachers**

0606 ADDITIONAL MATHEMATICS

0606/01

Paper 1, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2009 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2009	0606	01

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Accuracy mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol \surd implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2, 1, 0 means that the candidate can earn anything from 0 to 2.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2009	0606	01

The following abbreviations may be used in a mark scheme or used on the scripts:

AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through $\sqrt{}$ ” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy.
OW –1,2	This is deducted from A or B marks when essential working is omitted.
PA –1	This is deducted from A or B marks in the case of premature approximation.
S –1	Occasionally used for persistent slackness – usually discussed at a meeting.
EX –1	Applied to A or B marks when extra solutions are offered to a particular equation. Again, this is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2009	0606	01

<p>1 (i) $12 = 15\theta$, $\theta = 0.8$ rads</p> <p>(ii) Area = $\frac{1}{2}15^2(0.8)$ leading to 90 (cm²)</p>	<p>M1, A1 [2]</p> <p>M1</p> <p>A1</p> <p>[2]</p>	<p>M1 for use of $s = r\theta$</p> <p>M1 for use of $A = \frac{1}{2}r^2\theta$</p>
<p>2 $x^3 = 8$, leading to $x = 2$</p> <p>$\frac{dy}{dx} = 3x^2$ leading to grad of $-\frac{1}{12}$ for normal</p> <p>$y - 0 = -\frac{1}{12}(x - 2)$</p> <p>$\left(y = -\frac{1}{12}x + \frac{1}{6}\right)$</p>	<p>B1</p> <p>M1</p> <p>DM1</p> <p>A1</p> <p>[4]</p>	<p>B1 for finding where curve crosses the x axis</p> <p>M1 for attempt to differentiate and use of $m_1m_2 = -1$</p> <p>DM1 for attempt at equation of normal Allow unsimplified</p>
<p>3</p> <p>$\frac{1 - \cos^2 \theta}{\sec^2 \theta - 1} = \frac{\sin^2 \theta}{\tan^2 \theta}$ $= \cos^2 \theta$ $= 1 - \sin^2 \theta$</p> <p>Alt Scheme</p> <p>$\frac{1 - \cos^2 \theta}{\sec^2 \theta - 1} = \frac{\sin^2 \theta}{1 - \cos^2 \theta / \cos^2 \theta}$ $= \frac{\sin^2 \theta \cos^2 \theta}{\sin^2 \theta}$ $= \cos^2 \theta$ $= 1 - \sin^2 \theta$</p>	<p>M1</p> <p>M1</p> <p>M1</p> <p>A1 [4]</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p>	<p>M1 for use of $1 - \cos^2 \theta = \sin^2 \theta$</p> <p>M1 for use of $\sec^2 \theta - 1 = \tan^2 \theta$</p> <p>M1 for attempt to simplify</p> <p>M1 for use of $1 - \cos^2 \theta = \sin^2 \theta$</p> <p>M1 for attempting to get all in terms of cos</p> <p>M1 for attempt to simplify</p>
<p>4 (i) $5x - 3 = kx^2 - 3x + 5$ $kx^2 - 8x + 8 = 0$ using $b^2 - 4ac = 0$, $k = 2$ (Alt scheme: $5 = 2kx - 3$, $x = \frac{4}{k}$ $\frac{20}{k} - 3 = \frac{16}{k} - \frac{12}{k} + 5$ leading to $k = 2$)</p> <p>(ii) leading to $x = 2$, $y = 7$</p>	<p>M1</p> <p>DM1, A1 [3]</p> <p>M1, A1 [2]</p>	<p>M1 for equating line and curve equations</p> <p>DM1 for use of $b^2 - 4ac$ on resulting quadratic</p> <p>(Alt scheme: M1 for attempt to differentiate quadratic and equate to 5 DM1 for simplification and solution using resulting quadratic)</p> <p>M1 for obtaining x and y coords</p>

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2009	0606	01

<p>5 (a) $3^{2(2x-1)} = 3^{3x}$ $4x - 2 = 3x$ $x = 2$</p> <p>(b) $a^{-2}b$ or $\frac{b}{a^2}$ (allow here) $p = -2, q = 1$</p>	<p>B1 B1 B1 [3]</p> <p>B1 B1 [2]</p>	<p>B1 for $3^{2(2x-1)}$ B1 for 3^{3x} B1 for $x = 2$</p> <p>B1 for each</p>
<p>6 $f(3), f(-5)$ or $f(0.5) = 0$ spotted Either $(2x-1)(x^2+2x-15)$ Or $(x+5)(2x^2-7x+3)$ Or $(x-3)(2x^2+9x-5)$ $x = 3, -5, 0.5$</p>	<p>B1 M1 A1 M1 A2,1,0 [6]</p>	<p>B1 for spotting one root M1 for attempt to obtain quadratic factor A1 all correct M1 for solution of quadratic A2 for all 3 solutions (–1 each error) Correct factors only – lose 1 A mark</p>
<p>7 (i) $3xe^{3x} + e^{3x} - e^{3x}$ $= 3xe^{3x}$</p> <p>(ii) $\int xe^{3x} dx = \frac{1}{3} \left(xe^{3x} - \frac{e^{3x}}{3} \right)$</p>	<p>M1, A1, B1 [3]</p> <p>DM1 DM1 A1 [3]</p>	<p>M1 for attempt to differentiate a product. A1 for correct product. B1 for $-e^{3x}$</p> <p>DM1 for recognition of the ‘reverse’ to (i) DM1 for dealing with ‘3’ A1 all correct (condone omission of c)</p>
<p>8 (i) $\frac{dy}{dx} = \frac{(x^2+9)2 - 2x(2x)}{(x^2+9)^2}$ $= \frac{18-2x^2}{(x^2+9)^2}$, turning points, $x = \pm 3$</p> <p>(ii) $\frac{dx}{dt} = 2$ $\frac{dy}{dt} = 2 \times \left(\frac{16}{100} \right)$ $= 0.32$ or $\frac{8}{25}$</p>	<p>B2,1,0 M1 A1 [4]</p> <p>B1 M1 A1 [3]</p>	<p>Attempt to differentiate a quotient –1 each error</p> <p>M1 for correct attempt to find the turning points. A1 for both</p> <p>B1 for use of $\frac{dx}{dt} = 2$</p> <p>M1 for use of rates of change</p>

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2009	0606	01

<p>9 (i) $10\sqrt{2}\left(\frac{1}{\sqrt{2}}\mathbf{i} + \frac{1}{\sqrt{2}}\mathbf{j}\right) = 10\mathbf{i} + 10\mathbf{j}$</p> <p>(ii) $(-4\mathbf{i} + 8\mathbf{j}) + (20\mathbf{i} + 20\mathbf{j}) = 16\mathbf{i} + 28\mathbf{j}$</p> <p>(iii) $(10\mathbf{i} + 10\mathbf{j}) - (8\mathbf{i} + 6\mathbf{j}) = 2\mathbf{i} + 4\mathbf{j}$</p> <p>(iv) displacement of $(19\mathbf{i} + 34\mathbf{j}) - (16\mathbf{i} + 28\mathbf{j}) = 3\mathbf{i} + 6\mathbf{j}$ time = 1330 hours (accept 1.5 hours) at $3\mathbf{i} + 43\mathbf{j}$</p> <p>Alternative scheme: $(19\mathbf{i} + 34\mathbf{j}) + (8\mathbf{i} + 6\mathbf{j})t =$ $(16\mathbf{i} + 28\mathbf{j}) + (10\mathbf{i} + 10\mathbf{j})t$ or equivalent</p>	<p>M1 A1 [2]</p> <p>M1 A1 [2]</p> <p>M1 A1 [2]</p> <p>M1 A1 [3]</p>	<p>M1 for attempt at a correct direction vector A1 all correct</p> <p>M1 for valid attempt A1 all correct</p> <p>M1 for attempt at vector difference A1 condone negative</p> <p>M1 for displacement and attempt to obtain time A1 for correct time A1 for correct position vector</p> <p>M1 for attempt to equate like vectors A marks as above</p>
<p>10 (i) $m_{AB} = 0.75$ line AB $y - 0 = 0.75(x + 4)$ $m_{PQ} = -\frac{4}{3}$ line PQ $y - 10 = -\frac{4}{3}(x - 1)$ intersection at $C(4, 6)$ $Q(8.5, 0)$</p> <p>(ii) $AC = 10, CQ = 7.5$ Area = 37.5</p>	<p>M1 A1</p> <p>M1 A1 M1 A1 √ B1 [7]</p> <p>M1 A1 [2]</p>	<p>M1 for attempt at m_{AB} and line AB</p> <p>M1 for use of '$m_1m_2 = -1$' and attempt at line PQ</p> <p>M1 for attempt at solving simultaneous equations Ft on their line PQ</p> <p>M1 for attempt at lengths and area</p>

11 (i) $\ln s = n \ln t + \ln k$ <table border="1"> <tr> <td>$\ln t$</td><td>1.6</td><td>2.7</td><td>3.4</td><td>4.2</td><td>4.6</td></tr> <tr> <td>$\ln s$</td><td>7.2</td><td>5.9</td><td>5</td><td>4</td><td>3.6</td></tr> </table> <p>Plot $\ln s$ against $\ln t$</p>						$\ln t$	1.6	2.7	3.4	4.2	4.6	$\ln s$	7.2	5.9	5	4	3.6
$\ln t$	1.6	2.7	3.4	4.2	4.6												
$\ln s$	7.2	5.9	5	4	3.6												
(ii) grad $n = -1.2$ (-1.4 to -1.0) Intercept = $\ln k$, leading to $k = 7900 - 10\,000$																	
(iii) when $t = 50$, $\ln t = 4.4$ leading to $s = 80$ ($72 - 92$)																	
Alternative method (i) $\lg s = n \lg t + \lg k$ <table border="1"> <tr> <td>$\lg t$</td><td>0.7</td><td>1.2</td><td>1.5</td><td>1.8</td><td>2</td></tr> <tr> <td>$\lg s$</td><td>3.1</td><td>2.5</td><td>2.2</td><td>1.7</td><td>1.6</td></tr> </table>						$\lg t$	0.7	1.2	1.5	1.8	2	$\lg s$	3.1	2.5	2.2	1.7	1.6
$\lg t$	0.7	1.2	1.5	1.8	2												
$\lg s$	3.1	2.5	2.2	1.7	1.6												

M1, A1
M1
A1

[4]

M1, A1
M1, A1

[4]

M1
A1


[2]

M1 for attempt to take logs
A1 for correct form
M1 for attempt to plot correct graph
A1 for a reasonable straight line

M1 for use of grad = n
M1 for use of intercept = $\ln k$

M1 for attempt to obtain s


Same scheme applies



Page 8	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2009	0606	01

<p>12 EITHER</p> <p>(i) amplitude = 1</p> <p>(ii) period = 6π, 18.8</p> <p>(iii) $\sin\left(\frac{x}{3}\right) = \frac{1}{2}$, $x = \frac{\pi}{2}, \frac{5\pi}{2}$</p> <p>(iv) Area under curve</p> $\int_{\frac{\pi}{2}}^{\frac{5\pi}{2}} \left(1 + \sin \frac{x}{3}\right) dx = \left[x - 3 \cos \frac{x}{3} \right]_{\frac{\pi}{2}}^{\frac{5\pi}{2}}$ <p>leading to $2\pi + 3\sqrt{3}$</p> <p>Area of rectangle = $\left(\frac{5\pi}{2} - \frac{\pi}{2}\right) \times \frac{3}{2}$</p> <p>$= 3\pi$</p> <p>Shaded area = $3\sqrt{3} - \pi$ (2.05)</p> <p>Alternative solution: Shaded area</p> $\int_{\frac{\pi}{2}}^{\frac{5\pi}{2}} \left(\sin \frac{x}{3} - 0.5\right) dx = \left[-0.5x - 3 \cos \frac{x}{3}\right]_{\frac{\pi}{2}}^{\frac{5\pi}{2}}$	<p>B1 [1]</p> <p>B1 [1]</p> <p>M1 A1, A1 [3]</p> <p>M1 B1, B1</p> <p>DM1</p> <p>M1</p> <p>A1 [6]</p> <p>M1 M1 B1, B1 DM1, A1</p>	<p>M1 for attempt to solve correctly A1 for each (allow degrees here)</p> <p>M1 for attempt to integrate B1 for x, B1 for $-3 \cos \frac{x}{3}$</p> <p>DM1 for correct use of limits</p> <p>M1 for attempt at rectangle plus subtraction – must be working in radians</p> <p>M1 for subtraction (must be using radians) M1 for attempt to integrate B1 for $-0.5x$, B1 for $-3 \cos \frac{x}{3}$ DM1 for correct use of limits</p>
--	--	---

Page 9	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2009	0606	01

<p>OR</p> <p>(i) $t = \frac{\pi}{8}$</p> <p>(ii) $a = -4k \sin 4t$</p> <p>(iii) $12 = -4k \sin \frac{3\pi}{2}$ leading to $k = 3$</p> <p>(iv)</p>  <p>(v) $s = \int_0^{\frac{\pi}{24}} 3 \cos 4t \, dt$</p> <p>$= \left[\frac{3}{4} \sin 4t \right]_0^{\frac{\pi}{24}}$ leading to $\frac{3}{8}$</p>	<p>B1 [1]</p> <p>M1, A1 [2]</p> <p>M1 A1 [2]</p> <p>B1 √ B1 [2]</p> <p>M1, √ A1</p> <p>DM1, A1 [4]</p>	<p>M1 for attempt to differentiate</p> <p>M1 for attempt to substitute into their acceleration equation</p> <p>B1 for correct shape B1 ft on their value for k</p> <p>M1 for attempt to integrate Ft on their value for k</p> <p>DM1 for application of limits or equivalent</p>
--	--	--